


Distribution. Denmark: rare, and with decreasing frequency after 1950. In Jutland only in the southern and eastern parts. More frequent in F, LFM, SZ, NWZ and NEZ. No records from B. – Sweden: rather distributed in the south, north to Hls., but uncommon. Seemingly absent from southern Sk. and the western coastal districts (Hall-Boh.). – Norway: very rare, only SFi (1965) and MRi (1984). – Finland: fairly common in the southern and eastern districts, north to Sb and Kb; unrecorded from Al. Also in Vib and Kr of the USSR. – Central Europe; distribution not known in details.

Biology. In preference of habitat agreeing with, and sometimes occurring together with, *B. unipustulatus*. In moist, rather shaded sites, usually in mull-rich deciduous forest, for instance of ash, alder and beech, often in forest swamps. It dwells among moss and leaves. Mainly in May-June.



Figs 463-469. Penis of *Badister*. – 463: *bullatus* (Schränk); 464: *meridionalis* Puel; 465: *lacer-tosus* Sturm; 466: *sodalis* (Dft.); 467: *dilatatus* Chaud.; 468: *peltatus* (Pz.); 469: *anomalus* (Perris).

Figs 470, 471. Right paramere of *Badister*. – 470: *sodalis* (Dft.) and 471: *dorsiger* (Dft.).